

ON THE ORDER OF VANISHING OF THE CYCLOTOMIC p -ADIC L -FUNCTION

IVÁN BLANCO-CHACÓN

ABSTRACT. For a newform f for $\Gamma_0(N)$ of even weight k , we prove that the p -adic L -function $L_p(f, \alpha)$ is not identically zero on the group \mathbb{Z}_p^* of the p -adic units. If $p \geq 5$, we prove that the order of vanishing of $L_p(f, \alpha)$ at any p -adic integer is finite.

INTRODUCTION

To any eigenform $f \in S_k(\Gamma_0(N))$ of the Hecke operator T_p , Mazur-Tate-Teitelbaum [6] attached a p -adic L -function $L_p(f, \alpha; \chi)$ defined over $\mathcal{X} = \text{Hom}_{\text{cont}}(\mathbb{Z}_p^*, \mathbb{C}_p)$, the group of p -adic continuous characters. As a topological group, \mathcal{X} is isomorphic to $\mu_{p-1} \times \mathbb{Z}_p$, where μ_{p-1} stands for the group of $(p-1)$ -th roots of unity. From a theorem of Rohrlich ([8, Theorem 1]), it follows that if f is a normalized newform, $L_p(f, \alpha; \chi)$ does not vanish identically over \mathcal{X} , a conjecture stated for $k=2$ by Mazur and Swinnerton-Dyer ([5, Conjecture 1]).

The present article deals with the restriction of $L_p(f, \alpha; \chi)$ to the characters $\chi_s(x) = \langle x \rangle^s$, $s \in \mathbb{Z}_p^*$. We prove that the function $L_p(f, \alpha)(s) = L_p(f, \alpha; \chi_s)$ has a non-zero derivative at any p -adic integer s when f is a normalized newform. In particular, we obtain that $L_p(f, \alpha)$ does not vanish identically over $\{1\} \times \mathbb{Z}_p$, a refinement of Mazur and Swinnerton-Dyer's conjecture suggested by Henri Darmon.

In section 1, we review the construction of $L_p(f, \alpha; \chi)$, following [6]. In section 2, we apply an ultrametric analogue of the Stone-Weierstrass theorem to establish that $L_p(f, \alpha)$ does not vanish identically over \mathbb{Z}_p , provided that $p \geq 3$. If f has weight 2, we can say that, actually, $L_p(f, \alpha)$ does not vanish identically over \mathbb{Z}_p^* . In the ordinary case, the finiteness of the set of

Research funded by the FPI fellowship MTM2006-04895 and grant MTM2007-63582 of Spanish *Ministerio de Ciencia e Innovación*.

zeros of $L_p(f, \alpha)$ also follows, as well as the fact that if p is a primitive root mod N , and $\alpha \not\equiv 1 \pmod{p}$, this function does not vanish identically mod p , provided that not all the modular integrals are zero mod p .

In section 3, we use the above non-vanishing results in conjunction with some estimations of the moments of $L_p(f, \alpha)$ to prove our main result, namely, that the order of vanishing of $L_p(f, \alpha)(s)$ at any p -adic integer s is finite, if $p \geq 5$ (Theorem 3.1). In particular, it is finite at the central point $s = k/2$.

Acknowledgements. I want to thank my thesis advisor Pilar Bayer for her help and patience, and Henri Darmon for drawing my attention to this problem and suggesting some illuminating readings to begin with the resolution.

CONTENTS

Introduction	1
1. The cyclotomic p -adic L -functions	2
1.1. Modular integrals of cusp forms	2
1.2. The p -adic L -function of a cusp form	4
1.3. Special characters	6
2. Non-vanishing results	7
2.1. Approximation by polynomials	7
2.2. The non-vanishing on \mathbb{Z}_p	8
2.3. The non-vanishing mod p	11
3. Results on the order of the p -adic L -function	12
References	19

1. THE CYCLOTOMIC p -ADIC L -FUNCTIONS

1.1. Modular integrals of cusp forms. We fix p a prime number and embeddings $\overline{\mathbb{Q}} \hookrightarrow \mathbb{C}$, $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}_p}$. By \mathcal{O}_p , we denote the integral closure of \mathbb{Z} in $\overline{\mathbb{Q}_p}$. We consider the p -adic norm $|\cdot|$ on \mathbb{C}_p normalized so that $|p| = p^{-1}$.

The set of elements in \mathbb{Z}_p^* congruent to an integer $a \bmod p^n$, $p \nmid a$, will be denoted by $D(a, p^n)$.

Let $S_k(\Gamma_0(N))$ be the \mathbb{C} -vector space of cusp forms of positive even weight k for $\Gamma_0(N)$. Let \mathbb{T} be the \mathbb{Z} -algebra spanned by the Hecke operators $\{T_n\}_{n \geq 1}$ acting on it. In much of the paper, we will suppose that $f = \sum_{n \geq 1} a_n(f)q^n$ is a normalized newform. In this case, the number field $K_f = \mathbb{Q}(\{a_n(f)\})$ is totally real. Its ring of integers will be denoted by \mathcal{O}_f .

For the complex L -function $L(f, s)$, the following identity holds:

$$(1.1) \quad \Lambda(f, s) = N^{\frac{s}{2}}(2\pi)^{-s}\Gamma(s)L(f, s) = N^{\frac{s}{2}} \int_0^\infty f(it)t^{s-1}dt,$$

for $s \in \mathbb{C}$, $\operatorname{Re}(s) > \frac{k}{2} + 1$. The right hand side of the equality defines an entire function which satisfies the following functional equation:

$$(1.2) \quad \Lambda(f, s) = \pm \Lambda(f, k - s), \quad s \in \mathbb{C}.$$

If χ is a primitive Dirichlet character of conductor n , $(n, N) = 1$, and $\tau(\chi)$ its attached Gauss sum, the complex L -function of f twisted by χ is defined by

$$L(f, \chi, s) = \sum_{n=1}^{\infty} \frac{\chi(n)a_n(f)}{n^s}, \quad \operatorname{Re}(s) > \frac{k}{2} + 1.$$

Let $\Lambda(f, \chi, s) = N^{\frac{s}{2}}(2\pi)^{-s}\Gamma(s)L(f, \chi, s)$. By using orthogonality properties of the Dirichlet characters, we have that

$$(1.3) \quad \Lambda(f, \chi, s) = N^{\frac{s}{2}} \frac{\tau(\chi)}{n} \sum_{a=0}^{n-1} \overline{\chi(a)} \int_0^\infty f\left(\frac{a}{n} + it\right) t^{s-1} dt, \quad \operatorname{Re}(s) > \frac{k}{2} + 1.$$

The function $\Lambda(f, \chi, s)$ extends to an entire function and satisfies a functional equation similar to (1.2).

For $r \in \mathbb{Q}$ and $0 \leq j \leq k - 2$, the integrals

$$\lambda(f, r, j) = 2\pi \int_0^\infty f(r + it)t^j dt, \quad r \in \mathbb{Q},$$

are known in the literature as modular integrals. They satisfy the following

Theorem 1.1. ([6, § 2]) *There exists a \mathbb{Z} -lattice Σ_f of finite rank such that $\lambda(f, r, j) \in \Sigma_f$.*

It is an interesting question to study when the twisted L -function of a modular form vanishes. The following theorem is a result in this direction.

Theorem 1.2. (Rohrlich [8, Theorem 1]) *Let $f \in S_k(\Gamma_0(N))$ be a normalized newform. Let P be a finite set of primes and X_P the set of primitive Dirichlet characters unramified outside $P \cup \{\infty\}$. Then, for all but finitely many $\chi \in X_P$, $L\left(f, \chi, \frac{k}{2}\right) \neq 0$.*

Since any Dirichlet character χ of conductor p^n is unramified outside p , we deduce the following

Corollary 1.3. *For $n \in \mathbb{N}$ large enough there exists integers a_n , $p \nmid a_n$, such that*

$$\int_0^\infty f\left(\frac{a_n}{p^n} + it\right) t^{\frac{k}{2}-1} dt \neq 0.$$

1.2. The p -adic L -function of a cusp form. Let $f \in S_k(\Gamma_0(N))$ be an eigenform for T_p , $p \nmid N$, with eigenvalue $a_p(f)$. The Hecke polynomial attached to f at p is defined as follows:

$$X^2 - a_p(f)X + p^{k-1}.$$

For a root α of this polynomial and for $0 \leq j \leq k-2$, we may define an $\mathcal{O}_f[\frac{1}{\alpha}] \otimes \Sigma_f$ -valued p -adic distribution by setting

$$(1.4) \quad \mu_{\alpha,j}(D(a, p^n)) = \frac{1}{\alpha^n} \left(\lambda\left(f, \frac{a}{p^n}, j\right) - \frac{1}{\alpha} \lambda\left(f, \frac{a}{p^{n-1}}, j\right) \right), \quad n \geq 1.$$

If $|a_p(f)| = 1$, it is said that f is ordinary at p . Otherwise, f is said to be supersingular at p . A root α is admissible (cf. [6]) if $p^{1-k} < |\alpha| \leq 1$. In the ordinary case, there is only one admissible root, α ; it is a p -adic unit and $\mu_{\alpha,j}$ is then a p -adic measure. In the supersingular case, both roots are admissible, but the p -adic distributions are unbounded on the compact-open sets of \mathbb{Z}_p^* .

Definition 1.4. The cyclotomic distributions attached to f and p are defined as

- If f is ordinary at p : $\mu_{\alpha,j}$, where α is the unique root of the Hecke polynomial for f at p which is a p -adic unit.

- If f is supersingular at p : $\mu_{\alpha_i, j}$, $i = 1, 2$, where α_i denote the roots of the Hecke polynomial.

For an ordinary prime p , it is possible to integrate continuous \mathbb{Q}_p -valued functions by using uniform approximation by locally constant functions (cf. [4, Chapter 2]). The definition of an integral in the supersingular case requires some more work. First of all, we need to restrict the class of functions to be integrated.

Definition 1.5. A function $F : \mathbb{Z}_p^* \rightarrow \mathbb{Q}_p$ is said to be locally analytic if there is a covering of \mathbb{Z}_p^* by compact-open sets $D(a, p^m)$ such that

$$F|_{D(a, p^m)}(x) = \sum_{n=0}^{\infty} c_n(x-a)^n.$$

The following theorem provides an integral operator attached to any admissible root.

Theorem 1.6. (Mazur-Tate-Teitelbaum, [6]) *Let $f \in S_k(\Gamma_0(N))$ be an eigenform of the Hecke operator T_p . For a compact-open subset $K \subseteq \mathbb{Z}_p^*$, there exists a unique $\overline{\mathbb{Q}_p}^r$ -valued \mathbb{Q}_p -linear operator on the space of locally analytic functions, denoted by $\int_K F(x) d\mu_\alpha(x)$, with the following properties:*

- (1) *Interpolation property:* $\int_K x^j d\mu_\alpha(x) = \mu_{\alpha, j}(K)$, for $0 \leq j \leq k-2$.
- (2) *Divisibility property:* for any $n \geq 0$

$$\int_{D(a, p^n)} (x-a)^m d\mu_\alpha(x) \in \left(\frac{p^n}{\alpha}\right)^n \alpha^{-1} \Sigma_f \otimes \mathcal{O}_p.$$

- (3) *Continuity property:* if $F(x) = \sum_{n \geq 0} c_n(x-a)^n$ in $D(a, p^m)$, then

$$\int_{D(a, p^m)} F(x) d\mu_\alpha(x) = \sum_{n \geq 0} c_n \int_{D(a, p^m)} (x-a)^n d\mu_\alpha(x).$$

- (4) *For a fixed F , the assignment $K \mapsto \int_K F(x) d\mu_\alpha(x)$ yields a finitely additive function on the set of compact-open subsets of \mathbb{Z}_p^* .*

Let $\mathcal{X} = \text{Hom}_{\text{cont}}(\mathbb{Z}_p^*, \mathbb{C}_p)$. Since continuous characters are locally analytic, one can formulate the following

Definition 1.7. (The cyclotomic p -adic L -function) Assume that $p \nmid N$. Let $f \in S_k(\Gamma_0(N))$ be an eigenform of the Hecke operator T_p and α an admissible root of the Hecke polynomial of f at p . For $\chi \in \mathcal{X}$, we define

$$L_p(f, \alpha; \chi) = \int_{\mathbb{Z}_p^*} \chi(x) d\mu_\alpha(x).$$

1.3. Special characters. For $x \in \mathbb{Z}_p^*$, we write $x = \omega(x)\langle x \rangle$ where $\omega(x)$ is the unique $(p-1)$ -th root of unity in \mathbb{Z}_p^* congruent to $x \bmod p$ and $\langle x \rangle \in D(1, p)$.

Let us now consider the characters of the form $\chi_s(x) = \langle x \rangle^s$, for $s \in \mathbb{Z}_p$. For an admissible root α , we denote

$$L_p(f, \alpha)(s) = L_p(f, \alpha; \chi_{s-1}) = \int_{\mathbb{Z}_p^*} \langle x \rangle^{s-1} d\mu_\alpha(x), \quad s \in \mathbb{Z}_p.$$

As in the complex analytic, a functional equation holds for $L_p(f, \alpha)$. In order to recall it, we first introduce some notation.

Let $K \subseteq \mathbb{Z}_p^*$ be a compact-open set and F a locally analytic function over \mathbb{Z}_p^* . Let us define the map $g(x) = -\frac{1}{Nx}$, $x \in \mathbb{Z}_p^*$. We set

$$\tilde{F}(x) = N^{\frac{k-2}{2}} x^{k-2} (F \circ g), \quad \tilde{K} = g(K), \quad \tilde{f} = w_N(f),$$

where w_N stands for the Fricke involution on $S_k(\Gamma_0(N))$.

Proposition 1.8. (cf. [6, Theorem 17.1]) *If α is an admissible root for f ,*

$$\int_K F(x) d\mu_\alpha(x) = \int_{\tilde{K}} \tilde{F}(x) d\mu_\alpha(x).$$

Corollary 1.9. *Suppose that f is a newform. For any $s \in \mathbb{Z}_p$,*

$$L_p(f, \alpha)(s) = \pm \langle N \rangle^{1-s} \int_{\mathbb{Z}_p^*} x^{k-2} \langle x \rangle^{1-s} d\mu_\alpha(x).$$

In particular, if $k = 2$, $L_p(f, \alpha)(s) = 0$ if and only if $L_p(f, \alpha)(2-s) = 0$.

Proof. Since f is a normalized newform, it is an eigenfunction for the Fricke involution; hence, $\tilde{f} = \pm f$, and we have

$$L_p(f, \alpha)(s) = \int_{\mathbb{Z}_p^*} \langle x \rangle^{s-1} d\mu_\alpha(x) = \pm \int_{\mathbb{Z}_p^*} x^{k-2} \left\langle \frac{-1}{Nx} \right\rangle^{s-1} d\mu_\alpha(x).$$

Since $\langle \cdot \rangle$ is a multiplicative group homomorphism, the result follows. \square

2. NON-VANISHING RESULTS

If $p \nmid N$ and χ is a finite order p -adic character of conductor p^n , we have

$$(2.1) \quad L_p(\alpha, f; \chi\chi_j) = \frac{1}{\alpha^n} \left(1 - \frac{\overline{\chi(p)}p^{k-2}}{\alpha} \right) \left(1 - \frac{\chi(p)}{\alpha} \right) \frac{p^n}{\tau(\bar{\chi})} L(f, \bar{\chi}, j+1),$$

for $0 \leq j \leq k-2$ (cf. [6, pp. 20-21]). Rohrlich's theorem 1.2, together with identity 2.1, implies that $L_p(f, \alpha; \chi)$ does not vanish identically over \mathcal{X} . In this section we study the non-vanishing of $L_p(f, \alpha)$ over \mathbb{Z}_p .

2.1. Approximation by polynomials. We denote by χ_G the characteristic function of a set $G \subseteq \mathbb{Q}_p$. From now on, K will denote a compact-open subset of \mathbb{Q}_p . The norm of a continuous function $F : K \rightarrow \mathbb{Q}_p$ is defined by

$$\|F\|_{\infty, K} = \max_{x \in K} |F(x)|.$$

For an integer $n \geq 1$, let $K[X^n]$ denote the set of polynomials with coefficients in K spanned by the monomials $\{X^{nm}\}_{m \geq 0}$. Let $\overline{K[X^n]}$ be its closure in $\mathcal{C}(K, \mathbb{Q}_p)$ with respect to the ∞ -norm.

Definition 2.1. A function $F : K \rightarrow \mathbb{Q}_p$ is said to be a step function on K if there exists a finite family of compact-open subsets $G_i \subseteq K$, $1 \leq i \leq n$, such that $F = \sum_{i=1}^n a_i \chi_{G_i}$, with $a_i \in \mathbb{Q}_p$. A function $F : K \rightarrow \mathbb{Q}_p$ is said to be a locally polynomial function on K if for any $a \in K$ there exists a neighborhood G^a and a polynomial P_a such that $F|_{G^a}(s) = P_a(s)$, for any $s \in G^a$. If all the polynomials are of degree 0, we say that F is locally constant. We will denote by $\text{Loc}(K, \mathbb{Q}_p)$ the set of locally constant functions on K .

Proposition 2.2. *A function $F : K \rightarrow \mathbb{Q}_p$ is a locally polynomial function on K if and only if there exists a finite covering of K by compact-open subsets $\{G_i\}_{1 \leq i \leq n}$ and polynomials $\{P_i\}_{1 \leq i \leq n}$ with coefficients in \mathbb{Q}_p such that $F(x) = \sum_{i=1}^n P_i(x) \chi_{G_i}(x)$, for any $x \in K$. In particular, F is locally constant if and only if F is a step function.*

Theorem 2.3. ([1, Proposition 2.6]) *If A is a closed subalgebra of $\mathcal{C}^n(K, \mathbb{Q}_p)$ which separates points and contains the constant functions, then $\text{Loc}(K, \mathbb{Q}_p) \subseteq A$.*

Corollary 2.4. *If A is a closed subalgebra of $\mathcal{C}^n(K, \mathbb{Q}_p)$ that separates points and contains the constant functions, then A contains the locally polynomial functions on K with coefficients in \mathbb{Q}_p .*

Proof. Let $F = \sum_{i=1}^n P_i \chi_{G_i}$ be a locally polynomial function with coefficients in \mathbb{Q}_p . For any $\varepsilon > 0$, let us consider $\varepsilon_0 = \varepsilon / \max_{1 \leq i \leq n} \|P_i\|_{\infty, K}$. By 2.3, there exist $g_i \in A$ such that $\|g_i - \chi_{G_i}\|_{\infty, K} < \varepsilon_0$. The result now follows. \square

Corollary 2.5. $\text{Loc}(\mathbb{Z}_p^*, \mathbb{Q}_p) \subseteq \overline{\mathbb{Z}_p^*[X^{p-1}]}$.

Proof. Since the function $F(x) = x^{p-1}$ is injective on $U = D(1, p)$, it follows from Theorem 2.4 that $\text{Loc}(U, \mathbb{Q}_p) \subseteq \overline{U[X^{p-1}]}$, and since the balls $\{D(1, p^n)\}_{n \geq 1}$ form a basis of the topology of U , for any $n \geq 1$ we can find a sequence $(P_m(X))$ of polynomials in $U[X^{p-1}]$ such that

$$\lim_{m \rightarrow \infty} \max_{x \in D(1, p^n)} |P_m(x^{p-1}) - \chi_{D(1, p^n)}(x)| = 0.$$

But, given $1 \leq a \leq p^n - 1$ coprime to p , $\chi_{D(a, p^n)}(x) = \chi_{D(1, p^n)}(xa^{-1})$ and

$$\begin{aligned} & \max_{x \in D(1, p^n)} |P_m(x^{p-1}) - \chi_{D(1, p^n)}(x)| \\ &= \max_{ax \in D(a, p^n)} |P_m(a^{1-p}(xa)^{p-1}) - \chi_{D(1, p^n)}\left(\frac{ax}{a}\right)| \\ &= \max_{y \in D(a, p^n)} |Q_m(x^{p-1}) - \chi_{D(a, p^n)}(y)|. \end{aligned}$$

Since $Q_m(X^{p-1}) \in \mathbb{Z}_p^*[X^{p-1}]$ and the compact-open sets $\{D(a, p^n)\}$ form a covering of \mathbb{Z}_p^* , the result holds. \square

2.2. The non-vanishing on \mathbb{Z}_p . We will use the following

Lemma 2.6. *Assume that $p > 2$, let $f \in S_k(\Gamma_0(N))$ be a normalized newform and α an admissible root of the Hecke polynomial for f at p . If the p -adic integral operator attached to f , p and α vanishes over $\mathbb{Z}_p^*[X^{p-1}]$, then*

$$\int_{\mathbb{Z}_p^*} x^j \chi_{D(a, p^n)}(x) d\mu_\alpha(x) = 0,$$

for any $0 \leq j \leq k-2$, $n \geq 1$, and any a coprime to p , $1 \leq a \leq p^n - 1$.

Proof. a) If f is ordinary at p , the unique admissible root α is a p -adic unit and $\mu_{\alpha,j}$ are p -adic measures. Thus, if we set $F_j(x) = x^j \chi_{D(a,p^n)}(x)$, for $x \in \mathbb{Z}_p^*$, we may choose a subset of polynomials $\{P_m\} \subseteq \mathbb{Z}_p^*[X^{p-1}]$ such that $\lim_{m \rightarrow \infty} \|P_m - F_j\|_{\infty, \mathbb{Z}_p^*} = 0$. But $\int_{\mathbb{Z}_p^*} P_m(x) d\mu_\alpha(x) = 0$ for any $m \geq 1$, and since there exists $M > 0$ such that $|\mu_{\alpha,j}(K)| < M$ for any compact-open subset $K \subseteq \mathbb{Z}_p^*$, we have

$$\left| \int_{\mathbb{Z}_p^*} F_j(x) d\mu_\alpha(x) \right| = \left| \int_{\mathbb{Z}_p^*} (P_m(x) - F_j(x)) d\mu_\alpha(x) \right| \leq M \|P_m - F_j\|_{\infty, \mathbb{Z}_p^*},$$

which tends to zero when $m \rightarrow \infty$.

b) If f is supersingular at p , none of the distributions is bounded and we need to change the strategy. We choose a subset of polynomials $\{P_m\} \subseteq \mathbb{Z}_p^*[X^{p-1}]$ such that

$$\|P_m - F_j\|_{\infty, \mathbb{Z}_p^*} < p^{-rm}, \text{ for any } r \geq 1.$$

This means that

$$P_m(x) - F_j(x) = p^{rm} \sum_{j=r+1}^{N_m} a_m(x-a)^j, \text{ for any } x \in D(a, p^n),$$

with $a_m \in \mathbb{Z}_p$ and N_m the degree of P_m . On the other hand, since $P_m - F_j$ is a locally polynomial function on \mathbb{Z}_p^* with coefficients in \mathbb{Z}_p , if $b \neq a$, there exists $Q_{b,m}^a \in \mathbb{Q}_p[X]$ such that the restriction of $P_m - F_j$ to $D(b, p^n)$ equals $p^{rm} Q_{b,m}^a$, for any $x \in D(b, p^n)$. Thus $Q_{b,m}^a \in \mathbb{Z}_p[X]$. Now

$$\begin{aligned} \int_{D(a, p^n)} P_m d(x) \mu_\alpha(x) - \int_{D(a, p^n)} F_j(x) d\mu_\alpha(x) = \\ p^{rm} \sum_{k=r+1}^{N_m} a_m \int_{D(a, p^n)} (x-a)^k d\mu_\alpha(x). \end{aligned}$$

But $\int_{D(a, p^n)} (x-a)^k d\mu_\alpha(x) = \alpha^{-n-2} p^{nk} \omega_{k,m}$ with $\omega_{k,m} \in \mathcal{O}_p \otimes \Sigma_f$. Since Σ_f is an \mathcal{O}_f -lattice, all the $\omega_{k,m}$ are bounded. Thus, the right hand side tends to 0 p -adically when r increases. If $b \neq a$, $\int_{D(b, p^n)} F_j(x) d\mu_\alpha(x) = 0$ and

$$\int_{D(b, p^n)} P_m(x) d\mu_\alpha(x) - \int_{D(b, p^n)} F_j(x) d\mu_\alpha(x) = p^{rm} \int_{D(b, p^n)} Q_{b,m}^a(x) d\mu_\alpha(x).$$

Since n is fixed and the Taylor expansion of $Q_{b,m}^a$ around b has coefficients in \mathbb{Z}_p , all the integrals multiplying the p^{rm} term are bounded; thus the norm of the right hand side also tends to 0 when r increases. Therefore,

$$\int_{\mathbb{Z}_p^*} P_m(x) d\mu_\alpha(x) = 0 = \int_{\mathbb{Z}_p^*} F_j(x) d\mu_\alpha(x) + p^{rm} K_m,$$

with $|K_m|$ bounded, so that $\int_{\mathbb{Z}_p^*} F_j(x) d\mu_\alpha(x) = 0$. \square

The main result of this section is the following

Theorem 2.7. *Let $p > 2$ and let $f \in S_k(\Gamma_0(N))$ be a normalized newform. Then for any admissible root α of the Hecke polynomial of f at p , the p -adic function $L_p(f, \alpha)$ does not vanish identically over \mathbb{Z}_p . Furthermore, if $k = 2$, it does not vanish identically over \mathbb{Z}_p^* .*

Proof. Let us prove first that $L_p(f, \alpha)$ does not vanish identically over \mathbb{Z}_p . If this were not the case, we would have

$$\int_{\mathbb{Z}_p^*} \langle x \rangle^r d\mu_\alpha(x) = 0, \quad \text{for all } r \geq 0.$$

Thus $\int_{\mathbb{Z}_p^*} \langle x \rangle^{m(p-1)} d\mu_\alpha(x) = \int_{\mathbb{Z}_p^*} x^{m(p-1)} d\mu_\alpha(x) = 0$, for any $m \geq 0$. Now, we can apply Lemma 2.6 to conclude that, in particular,

$$\int_{D(a, p^n)} x^{\frac{k}{2}-1} d\mu_\alpha(x) = 0.$$

But this would imply that for any $n \geq 1$, a coprime to p , and $1 \leq a \leq p^n - 1$,

$$(2.2) \quad \int_0^\infty f\left(\frac{a}{p^n} + it\right) t^{\frac{k}{2}-1} dt - \frac{1}{\alpha} \int_0^\infty f\left(\frac{a}{p^{n-1}} + it\right) t^{\frac{k}{2}-1} dt = 0.$$

Since f is a cusp form, setting $q = e^{2\pi iz}$, $\text{Im}(z) > 0$, there exist constants $M, C_f > 0$ such that

$$\left| \frac{f(q)}{q} \right| \leq C_f, \quad \text{if } |q| < M.$$

Thus if $y \gg 1$, $|f(x + iy)| < C_f e^{-2\pi y}$, and for any a coprime to p ,

$$(2.3) \quad \lim_{m \rightarrow \infty} \int_0^\infty f\left(\frac{a}{p^m} + it\right) t^{\frac{k}{2}-1} dt = \int_0^\infty f(it) t^{\frac{k}{2}-1} dt.$$

If $\alpha \neq 1$, by taking limits in 2.2, and taking into account 2.3 and the fact that f is a periodic function of period 1, we would have

$$\int_0^\infty f(a+it)t^{\frac{k}{2}-1}dt = 0, \quad \text{for any } a \in \mathbb{Z}.$$

Thus, applying 2.2 recursively, we would obtain

$$\int_0^\infty f\left(\frac{a}{p^m} + it\right)t^{\frac{k}{2}-1}dt = 0,$$

which contradicts Corollary 1.3.

If $\alpha = 1$, setting $n = 1$ in 2.2, we would have

$$\int_0^\infty f\left(\frac{a}{p} + it\right)t^{\frac{k}{2}-1}dt = \int_0^\infty f(it)t^{\frac{k}{2}-1}dt,$$

and, replacing it recursively in 2.2, we would obtain

$$\int_0^\infty f\left(\frac{a}{p^m} + it\right)t^{\frac{k}{2}-1}dt = \int_0^\infty f(it)t^{\frac{k}{2}-1}dt,$$

for any a coprime to p . But this would yield

$$\Lambda\left(f, \chi, \frac{k}{2}\right) = N^{\frac{s}{2}} \frac{\tau(\chi)}{p^n} \sum_{a=1}^{p^n-1} \overline{\chi(a)} \int_0^\infty f(it)t^{\frac{k}{2}-1}dt = 0,$$

for any Dirichlet character χ , which contradicts Theorem 1.2.

If, moreover, $k = 2$, the result follows from the functional equation in Corollary 1.9, because, if $|s| < 1$, then $2 - s$ is p -adic unit. \square

When $k = 2$, it is particularly interesting to restrict the domain of $L_p(f, \alpha; \chi)$ to \mathbb{Z}_p , since it yields a natural way of associating p -adic L -functions to modular abelian varieties.

2.3. The non-vanishing mod p . We say that a modular integral $\lambda \in \Sigma_f$ is divisible by p if its coordinates are valued in $D(0, p)^r$, r being the rank of Σ_f .

Proposition 2.8. *Let $p > 2$ be an ordinary prime for $f \in S_k(\Gamma_0(N))$, and assume that p is a primitive root mod N . Suppose that $\alpha \not\equiv 1 \pmod{p}$. If $L_p(f, \alpha)$ vanishes identically mod p over \mathbb{Z}_p^* , then all the modular integrals are divisible by p .*

Proof. Let us first observe that if all the modular integrals $\lambda(f, \frac{a}{p^n}, j)$ are divisible by p , then $\lambda(f, 0, j)$ is also divisible by p , since $\lim_{n \rightarrow \infty} \lambda(f, \frac{a}{p^n}, j) = \lambda(f, 0, j)$ and Σ_f is closed in \mathbb{C}^r . Thus, we are reduced to proving that, under the hypothesis of the proposition, the modular integrals $\lambda(f, \frac{a}{p^m}, j)$, with a coprime to p and $0 \leq j \leq k-2$ are divisible by p . Assuming that this were not the case, there would exist j_0 , a minimal $n_0 \geq 1$, and an integer a_0 coprime to p such that $p \nmid \lambda(f, \frac{a_0}{p^{n_0}}, j_0)$. Hence, the measure $\mu_{p,\alpha,j_0}(D(a_0, p^{n_0}))$ would not be divisible by p . On the other hand, by Corollary 2.4, there would exist a sequence $\{P_k(X)\}$ of polynomials in $\mathbb{Z}_p[X^{p-1}]$, converging to the locally polynomial function $x^{j_0} \chi_{D(a_0, p^{n_0})}(x)$ such that

$$\lim_{k \rightarrow \infty} \int_{\mathbb{Z}_p^*} P_k(x) d\mu_\alpha(x) = \mu_{p,\alpha,j_0}(D(a_0, p^{n_0})).$$

Since, for any k , $\int_{\mathbb{Z}_p^*} P_k(x) d\mu_\alpha(x) \in D(0, p)^r$, and this is a closed subset, we would have $\mu_{p,\alpha,j_0}(D(a_0, p^{n_0})) \in D(0, p)^r$, which is a contradiction. To complete the proof, let us notice that, p being a primitive root mod N , any modular symbol is a linear combination with integer coefficients of the modular symbols $\lambda\left(f, \frac{a_{k_i}}{p^{n_{k_i}}}, j_i\right)$.

□

3. RESULTS ON THE ORDER OF THE p -ADIC L-FUNCTION

We are going to exploit the fact that the p -adic L -function $L_p(f, \alpha)$ is not identically zero on \mathbb{Z}_p to prove our main result.

Theorem 3.1. *Let $f \in S_k(\Gamma_0(N))$ be a normalized newform, $p \geq 5$, and α an admissible root of the Hecke polynomial for f at p . Then, for any $s_0 \in \mathbb{Z}_p$,*

$$\text{ord}_{s=s_0} L_p(f, \alpha)(s) < \infty.$$

We recall the following standard notation

$$l_\infty(\mathbb{C}_p) = \{(x_n)_{n \geq 1} \in \mathbb{C}_p^{\mathbb{N}} : \sup_{n \geq 1} |x_n| < \infty\},$$

$$c_0(\mathbb{C}_p) = \{(x_n)_{n \geq 1} \in \mathbb{C}_p^{\mathbb{N}} : \lim_{n \rightarrow \infty} |x_n| = 0\}.$$

Both linear spaces are complete with respect to the norm

$$\|(x_n)_{n \geq 1}\|_\infty = \sup_{n \geq 1} |x_n|.$$

Let $D_i = D(i, p)$, $1 \leq i \leq p-1$. Given $x \in D_i$ and by setting ω_i for the unique $(p-1)$ -th root of unity congruent to $i \pmod p$, we may write $x = \omega_i \langle x \rangle$, with $\langle x \rangle = 1 + p\tilde{x}$. Then

$$\tilde{x} = \frac{1}{p} \left(\frac{x}{\omega_i} - 1 \right) = \frac{1}{p} \left(\frac{1}{\omega_i} (x - i) + \frac{i}{\omega_i} - 1 \right).$$

We set $a_{1,i} = p^{-1}\omega_i^{-1} \in p^{-1}\mathbb{Z}_p$ and $a_{0,i} = p^{-1}(i\omega_i^{-1} - 1) \in \mathbb{Z}_p$.

For an admissible root α , let us write

$$L_p(f, \alpha)(s) = \int_{\mathbb{Z}_p^*} (1 + p\tilde{x})^{s-1} d\mu_\alpha(x) = \int_{\mathbb{Z}_p^*} \sum_{k=0}^{\infty} \binom{s-1}{k} p^k \tilde{x}^k d\mu_\alpha(x).$$

Since $n! = p^{\frac{n-\sigma_n}{p-1}} u_n$, with $u_n \in \mathbb{Z}_p^*$ and σ_n being equal to the sum of the p -adic digits of n , we shall have

$$\binom{s-1}{k} p^k = p^{\frac{(p-2)k+\sigma_k}{p-1}} u_k q_k(s),$$

where $q_k(s) = (s-1)(s-2)\dots(s-k) \in \mathbb{Z}_p[s]$ is a polynomial of degree k .

Proposition 3.2. *If $p \geq 5$, for any sequence $(u_k)_{k \geq 1}$ of p -adic units, the sequence*

$$\left(p^{\frac{(p-2)k+\sigma_k}{p-1}} u_k \int_{\mathbb{Z}_p^*} \tilde{x}^k d\mu_\alpha(x) \right)_{k \geq 1} \text{ is in } c_0(\mathbb{C}_p).$$

Proof. Let $g^k(x) = \tilde{x}^k$, $k \geq 1$. With the former notations, in each disc D_i we shall have

$$g|_{D_i}(x) = a_{1,i}(x - i) + a_{0,i},$$

with $a_{1,i} \in p^{-1}\mathbb{Z}_p$ and $a_{0,i} \in \mathbb{Z}_p$. Thus,

$$g_k|_{D(i)}(x) = \sum_{n=0}^k \binom{k}{n} a_{1,i}^n a_{0,i}^{k-n} (x - i)^n.$$

Furthermore, we may write

$$\int_{D(i,p)} (x - i)^n d\mu_\alpha(x) = \alpha^{-2} p^n \gamma_{i,n}, \text{ with } \gamma_{i,n} \in \Sigma_f \otimes \mathcal{O}_p.$$

Thus,

$$\left| \int_{D(i,p)} g_k(x) d\mu_\alpha(x) \right| \leq R |\alpha^{-2}| \max_{0 \leq n \leq k} \left| \binom{k}{n} \right| \leq R |\alpha^{-2}| p^{\frac{k-\sigma_k}{p-1}}.$$

Combining this with the strong triangle inequality, we obtain

$$\left| \int_{\mathbb{Z}_p^*} \tilde{x}^k d\mu_\alpha(x) \right| \leq R |\alpha^{-2}| p^{\frac{k-\sigma_k}{p-1}},$$

so that

$$|x_k| = p^{\frac{(2-p)k-\sigma_k}{p-1}} \left| \int_{\mathbb{Z}_p^*} \tilde{x}^k d\mu_\alpha(x) \right| \leq R |\alpha^{-2}| p^{\frac{(3-p)k-\sigma_k}{p-1}}.$$

□

Using Proposition 3.2 together with the fact that $\frac{p^k}{k!} = p^{\frac{(p-2)k+\sigma_k}{p-1}} u_k$, with $u_k \in \mathbb{Z}_p^*$, we can ensure that the series

$$\langle x \rangle^{s-1} = \sum_{k=0}^{\infty} \binom{s-1}{k} p^k \tilde{x}^k,$$

converges uniformly in \mathbb{Z}_p^* . Thus,

$$(3.1) \quad L_p(f, \alpha)(s) = \sum_{k=0}^{\infty} \int_{\mathbb{Z}_p^*} \binom{s-1}{k} p^k \tilde{x}^k d\mu_\alpha(x).$$

The following result will allow us to differentiate the series $L_p(f, \alpha)(s)$ term by term.

Proposition 3.3. *Let $f_n(s) = n! \binom{s-1}{n}$. If $f(s) = \sum_{n=0}^{\infty} \lambda_n f_n(s)$ with $(\lambda_n)_{n \geq 0} \in c_0(\mathbb{C}_p)$, then the series converges uniformly in \mathbb{Z}_p and*

$$f^{(j)}(s) = \sum_{n=0}^{\infty} \lambda_n f_n^{(j)}(s), \quad \text{for } j \geq 0.$$

Proof. Let us first observe that, since $(\lambda_n)_n \in c_0(\mathbb{C}_p)$, the sum $\sum_{n=0}^{\infty} \lambda_n f^{(j)}(s)$ converges for any $s \in \mathbb{Z}_p^*$. We have to prove that, for any $j \geq 0$,

$$(3.2) \quad \lim_{m \rightarrow \infty} \left| \frac{\sum_{n=0}^{\infty} \lambda_n \left(f_n^{(j)}(s + p^m) - f_n^{(j)}(s) - p^m f_n^{(j+1)}(s) \right)}{p^m} \right| = 0, \quad \text{for any } s \in \mathbb{Z}_p^*.$$

We remark that the mean-value theorem does not hold in general in the p -adic setting (cf. [7]). We now proceed by induction on j .

a) For $j = 1$, Let us denote by $e_k(a_1, \dots, a_n)$ the k -th symmetrical elementary polynomial in the roots $\{a_1, \dots, a_n\}$. For $n, m \geq 1$ and $x \in \mathbb{Z}_p^*$, we have that

$$f_n(s) = e_n(s-1, \dots, s-n),$$

$$f_n(s+p^m) = \sum_{j=0}^n p^{jm} e_{n-j}(s-1, \dots, s-n),$$

and $f'_n(s) = e_{n-1}(s-1, s-2, \dots, s-n)$. Thus,

$$f_n(s+p^m) - f_n(s) - p^m f'_n(s) = \sum_{j=2}^n p^{jm} e_{n-j}(s-1, \dots, s-n)$$

and

$$\left| \frac{f_n(s+p^m) - f_n(s) - p^m f'_n(s)}{p^m} \right| \leq p^{-m}.$$

Since $(\lambda_n)_n$ is bounded, the result follows in this case.

b) Assume that the result holds for $j \geq 1$. A straightforward computation shows that

$$f_n^{(j)}(s) = j! e_{n-j}(s-1, s-2, \dots, s-n).$$

Thus,

$$f_n^{(j)}(s+p^m) = j! \sum_{k=0}^{n-j} c_k p^{km} e_{n-j-k}(s-1, s-2, \dots, s-n),$$

for suitable integers $c_k \geq 1$.

Lemma 3.4. $c_1 = j+1$. □

Let us note that

$$f_n^{(j+1)}(s) = (j+1)! e_{n-j-1}(s-1, s-2, \dots, s-n).$$

Hence, using Lemma 3.4, we have

$$f_n^{(j)}(s+p^m) - f_n^{(j)}(s) - p^m f_n^{(j+1)}(s) = j! \sum_{k=2}^{n-j} c_k p^{km} e_{n-j-k}(s-1, s-2, \dots, s-n)$$

and the result holds. □

We proceed now with the proof of Theorem 3.1.

Proof. Fix $s_0 \in \mathbb{Z}_p$. Since $L_p(f, \alpha)$ does not vanish identically, there exists an integer $k \geq 0$ such that

$$\int_{\mathbb{Z}_p^*} \tilde{x}^k d\mu_\alpha(x) \neq 0.$$

Let us define the (non-empty) set

$$\Sigma = \{k \geq 0 : \int_{\mathbb{Z}_p^*} \tilde{x}^k d\mu_{f,\alpha} \neq 0\}.$$

If Σ is finite, we consider its maximal integer k and, by taking the k -th derivative of $L_p(f, \alpha)(s)$ at $s = s_0$, we obtain

$$\frac{d^k}{ds^k} L_p(f, \alpha)(s)|_{s=s_0} = p^k \int_{\mathbb{Z}_p^*} \tilde{x}^k d\mu_{f,\alpha} \neq 0.$$

Now we suppose that Σ is an infinite set. The k_n -th term in the series has the form

$$p^{k_n} \binom{s-1}{k_n} \int_{\mathbb{Z}_p^*} \tilde{x}^{k_n} d\mu_{f,\alpha} = p^{\frac{(p-2)k_n + \sigma_{k_n}}{p-1}} u_{k_n} q_{k_n}(s) \int_{\mathbb{Z}_p^*} \tilde{x}^{k_n} d\mu_{f,\alpha},$$

with $q_{k_n}(s) = (s-1)(s-2)\dots(s-k_n)$. If we denote by $a^{i,j}$ the k_i -th derivative of $q_{k_j}(s)$ at $s = s_0$, we may define the linear endomorphism

$$\psi : c_0(\mathbb{C}_p) \longrightarrow c_0(\mathbb{C}_p), \quad x = (x_n) \mapsto \sum_{r=n}^{\infty} a^{r,n} x_n.$$

Since $a^{i,j} \in \mathbb{Z}_p$, ψ is well defined and $\|\psi(x)\| \leq \|x\|$, it is a continuous map. We can see this endomorphism as being represented by an infinite matrix $(a^{i,j})_{i,j \geq 1}$ which is upper triangular.

Proposition 3.5. *The endomorphism ψ is injective.*

Proof. For $x = (x_n)_{n \geq 1} \in c_0(\mathbb{C}_p)$, we have $\psi(x) = (D \circ \phi)(x)$ where $D : c_0(\mathbb{C}_p) \longrightarrow c_0(\mathbb{C}_p)$, $x = (x_n) \mapsto (k_n! x_n)$, and $\phi : c_0(\mathbb{C}_p) \longrightarrow c_0(\mathbb{C}_p)$ is given

by left natural multiplication with the infinite matrix

$$M_\phi = \begin{pmatrix} 1 & \frac{a_{1,2}}{k_1!} & \frac{a_{1,3}}{k_1!} & \frac{a_{1,4}}{k_1!} & \cdots \\ 0 & 1 & \frac{a_{2,3}}{k_2!} & \frac{a_{2,4}}{k_2!} & \cdots \\ 0 & 0 & 1 & \frac{a_{3,4}}{k_3!} & \cdots \\ 0 & 0 & 0 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Let us check that $\phi(c_0(\mathbb{C}_p)) \subseteq c_0(\mathbb{C}_p)$. Note that $q_{k_j}(s) = s^{k_j} + e_1^{k_j} s^{k_j-1} + \dots + e_{k_j}^{k_j}$, with $e_r^{k_j}$ the r -th elementary symmetrical polynomial in the roots $\{1, 2, \dots, k_j\}$. Thus,

$$\frac{1}{k_i!} \frac{d^{k_i}}{ds^{k_i}} q_{k_j}(s) = \binom{k_j}{k_j-k_i} s^{k_j-k_i} + \binom{k_j-1}{k_j-1-k_i} e_1^{k_j} s^{k_j-k_i-1} + \dots + \binom{k_i}{0} e_{k_i}^{k_j}.$$

Since

$$\binom{k_j-r}{k_j-k_i-r} = p^{\frac{-\sigma_{k_j-r} + \sigma_{k_j-k_i-r} + \sigma_{k_i}}{p-1}} \theta_r,$$

with $\theta_r \in \mathbb{Z}_p^*$, for $1 \leq r \leq k_j - k_i$, and

$$-\sigma_{k_j-r} + \sigma_{k_j-k_i-r} + \sigma_{k_i} = (p-1) \text{ord}_p \binom{k_j+k_i-r}{k_i}$$

(cf. [4]), it follows that

$$\left| \frac{1}{k_i!} \frac{d^{k_i}}{ds^{k_i}} q_{k_j}(s) \right|_{s=s_0} \leq 1.$$

Hence, ϕ is well defined and its associated matrix M_ϕ is upper triangular with ones in the diagonal. To prove the injectivity, we proceed by steps:

Step 1. We consider the adjoint matrix of the element $a_{i,j}$:

$$M_\phi^{i,j} = \begin{pmatrix} \frac{a_{1,1}}{k_1!} & \cdots & \frac{a_{1,j-1}}{k_1!} & \frac{a_{1,j+1}}{k_1!} & \cdots \\ 0 & \cdots & \frac{a_{2,j-1}}{k_2!} & \frac{a_{2,j+1}}{k_2!} & \cdots \\ \vdots & \cdots & \vdots & \vdots & \vdots \\ 0 & \cdots & \frac{a_{i-1,j-1}}{k_{i-1}!} & \frac{a_{i-1,j+1}}{k_{i-1}!} & \cdots \\ 0 & \cdots & \frac{a_{i+1,j-1}}{k_{i+1}!} & \frac{a_{i+1,j+1}}{k_{i+1}!} & \cdots \\ \vdots & \cdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

From a particular row onwards, this matrix becomes upper triangular, with ones on the diagonal, hence we can formally compute the determinant

$\det(M_\phi^{i,j})$ expanding along the first row. The result is a finite sum of finite products of the terms $\frac{a_{i,j}}{k_i!} \in \mathbb{Z}_p$, therefore, it is a p -adic integer.

Step 2. We put together all the adjoint matrices and take the transpose, obtaining an upper-triangular matrix with entries in \mathbb{Z}_p and ones on the diagonal. We write a few terms of this matrix:

$$M'_\phi = \begin{pmatrix} 1 & -\frac{a_{1,2}}{k_1!} & -\frac{a_{1,3}}{k_1!} + \frac{a_{1,2}a_{2,3}}{k_1!k_2!} & -\frac{a_{1,2}a_{2,3}a_{3,4}}{k_1!k_2!k_3!} + \frac{a_{1,3}a_{3,4}}{k_1!k_3!} + \frac{a_{1,2}a_{3,4}}{k_1!k_3!} - \frac{a_{1,4}}{k_1!} & \cdots \\ 0 & 1 & -\frac{a_{2,3}}{k_2!} & \frac{a_{2,4}a_{3,4}}{k_2!k_3!} - \frac{a_{2,4}}{k_2!} & \cdots \\ 0 & 0 & 1 & -\frac{a_{3,4}}{k_3!} & \cdots \\ 0 & 0 & 0 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

Now, M'_ϕ defines again a continuous linear endomorphism of $c_0(\mathbb{C}_p)$, that we call ϕ' .

Step 3. Since M_ϕ, M'_ϕ are upper triangular infinite matrices with ones on the diagonal and entries in \mathbb{Z}_p , we have that $(\phi' \circ \phi)(x) = x$, for all $x \in c_0(\mathbb{C}_p)$; i. e., ϕ' is a right inverse of ϕ .

Since D is trivially injective, so is ψ , completing the proof of Proposition 3.5 and Theorem 3.1. \square

To any elliptic curve E/\mathbb{Q} of conductor N , we can attach its complex analytic L -series $L(E, s)$. By modularity, there exists a weight 2 normalized newform f for $\Gamma_0(N)$ such that $L(E, s) = L(f_E, s)$. For an admissible root α of the Hecke polynomial of f_E at p , the corresponding p -adic L -function of E is defined by setting

$$L_p(E, \alpha)(s) = L_p(f_E, \alpha)(s), \quad s \in \mathbb{Z}_p.$$

Corollary 3.6. *If $p \geq 5$ and α is an admissible root of the Hecke polynomial of f_E at p , then*

$$\text{ord}_{s=1} L_p(E, \alpha)(s) < \infty.$$

REFERENCES

- [1] Araujo, J.; Schikhof, Wim H.: The Weierstrass-Stone approximation theorem for p -adic C^m -functions. *Ann. Math. Blaise Pascal* 1 (1994), no. 1, 61–74.
- [2] Darmon, H.: *Rational points on modular elliptic curves*. CBMS Regional Conference Series in Mathematics, 101. 2004.
- [3] Diamond, F.; Shurman, J.: *A first course in modular forms*. Graduate Texts in Mathematics, 228. Springer-Verlag, New York, 2005.
- [4] Koblitz, N.: *p -adic numbers, p -adic analysis, and zeta-functions*. Second edition. Graduate Texts in Mathematics, 58. Springer-Verlag, New York, 1984.
- [5] Mazur, B. ; Swinnerton-Dyer, P.: Arithmetic of Weil curves. *Invent. Math.* 25 (1974), 1–61.
- [6] Mazur, B.; Tate, J.; Teitelbaum, J.: On p -adic analogues of the conjectures of Birch and Swinnerton-Dyer. *Invent. Math.* 84 (1986), no. 1, 1–48.
- [7] Robert, Alain M.: *A course in p -adic analysis*. Graduate Texts in Mathematics, 198. Springer-Verlag, New York, 2000.
- [8] Rohrlich, David E.: L -functions and division towers. *Math. Ann.* 281 (1988), 611–632.

FACULTAD DE MATEMÁTICAS, UNIVERSIDAD DE BARCELONA, GRAN VIA DE LES CORTS CATALANES, 585. 08007 BARCELONA, SPAIN.